arXiv:1606.02787v1 [math.FA] 9 Jun 2016

Equivalent norms for the Morrey spaces with non-doubling measures

June 10, 2016

Yoshihiro Sawano, Hitoshi Tanaka[1]
Graduate School of Mathematical Sciences, The University of Tokyo,
3-8-1 Komaba, Meguro-ku Tokyo 153-8914, JAPAN
E-mail: yosihiro@ms.u-tokyo.ac.jp, htanaka@ms.u-tokyo.ac.jp

Abstract

In this paper under some growth condition we investigate the connection between RBMO and the Morrey spaces. We do not assume the doubling condition which has been a key property of harmonic analysis. We also obtain another type of equivalent norms.

KEYWORDS : Morrey space, Campanato space, equivalent norms

AMS Subject Classification : Primary 42B35, Secondary 46E35.

1 Introduction

In this paper we discuss equivalent norms for the (vector-valued) Morrey spaces with non-doubling measures. We consider the connection between the Morrey spaces and the Campanato spaces with underlying measure μ non-doubling. The Morrey spaces appeared in [5] originally in connection with the partial differential equations and the Campanato spaces in [1] and [2]. We refer to [6] for

[1] The first author is supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists. The second author is supported by the 21st century COE program at Graduate School of Mathematical Sciences, the University of Tokyo and by Fūjyukai foundation.

the result of Morrey spaces coming with the doubling measures. Before we state our main theorem, let us make a brief view of the terminology of measures on $\mathbf{R}^d$. We say that a (positive) Radon measure μ on $\mathbf{R}^d$ satisfies the growth condition if

$$\mu(Q(x,l)) \le C_0\, l^n \text{ for all } x \in \operatorname{supp}(\mu) \text{ and } l > 0, \tag{1}$$

where C_0 and $n \in (0, d]$ are some fixed numbers. A measure μ is said to satisfy the doubling condition if

$$\mu(Q(x,2l)) \le C\,\mu(Q(x,l)) \text{ for all } x \in \mathbf{R}^d \text{ and } l > 0$$

for some constant $C > 0$. A measure μ which satisfies the growth condition will be called growth measure while a measure μ with the doubling condition will be called the doubling measure.

By a "cube" $Q \subset \mathbf{R}^d$ we mean a closed cube having sides parallel to the axes. Its center will be denoted by z_Q and its side length by $\ell(Q)$. By $Q(x,l)$ we will also denote the cube centered at x of sidelength l. For $\rho > 0$, $\rho\, Q$ means a cube concentric to Q with its sidelength $\rho\,\ell(Q)$. Let $\mathcal{Q}(\mu)$ denote the set of all cubes $Q \subset \mathbf{R}^d$ with positive μ-measures. If μ is finite, we include $\mathbf{R}^d$ in $\mathcal{Q}(\mu)$ as well. In [7], the authors defined the Morrey spaces $\mathcal{M}^p_q(k,\mu)$ for non-doubling measures normed by

$$\|f : \mathcal{M}^p_q(k,\mu)\| := \sup_{Q\in\mathcal{Q}(\mu)} \mu(k\,Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f|^q\, d\mu\right)^{\frac{1}{q}},\ 1 \le q \le p < \infty,\ k > 1.$$

The fundamental property of this norm is

$$\|f : \mathcal{M}^p_q(k_1,\mu)\| \le \|f : \mathcal{M}^p_q(k_2,\mu)\| \le C_d \left(\frac{k_1-1}{k_2-1}\right)^d \|f : \mathcal{M}^p_q(k_1,\mu)\|$$

for $1 < k_1 < k_2 < \infty$. With this relation in mind, we will denote $\mathcal{M}^p_q(\mu) = \mathcal{M}^p_q(2,\mu)$. The aim of this paper is to find some norms equivalent to this Morrey norm.

2 Equivalent norm of doubling type

In this section we investigate an equivalent norm related to the doubling cubes. Although we now envisage the non-homogeneous setting, we are still able to place ourselves in the setting of the doubling cubes. In [10], Tolsa defined the notion of doubling cubes. Let $k, \beta > 1$. We say that $Q \in \mathcal{Q}(\mu)$ is a (k,β)-doubling cube, if $\mu(kQ) \le \beta\,\mu(Q)$. It is well-known that, if $\beta > k^d$, then for μ-almost all $x \in \mathbf{R}^d$ and for all $Q \in \mathcal{Q}(\mu)$ centered at x, we can find a (k,β)-doubling

cube from $k^{-1}Q, k^{-2}Q, \ldots$. In what follows we denote by $\mathcal{Q}(\mu; k, \beta)$ the set of all (k, β)-doubling cubes in $\mathcal{Q}(\mu)$. We fix $k, \beta > 1$ with $\beta > k^d$. Let $1 \le q \le p < \infty$. For $f \in L^1_{loc}(\mu)$ define

$$\|f \,:\, \mathcal{M}^p_q(\mu)\|_d := \sup_{Q \in \mathcal{Q}(\mu;k,\beta)} \mu(Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}}.$$

Now we present the main theorem in this section.

Theorem 2.1. *Let μ be a Radon measure which does not necessarily satisfy the growth condition nor the doubling condition and let $1 \le q < p < \infty$. If $\beta > k^{\frac{dpq}{p-q}}$, then*

$$C^{-1} \|f \,:\, \mathcal{M}^p_q(\mu)\|_d \le \|f \,:\, \mathcal{M}^p_q(\mu)\| \le C \|f \,:\, \mathcal{M}^p_q(\mu)\|_d, \ f \in \mathcal{M}^p_q(\mu),$$

for some constant $C > 0$.

Before we come to the proof of Theorem 2.1, two clarifying remarks may be in order.

Remark 2.2. If $p = q$, this theorem fails in general. However, if we assume the growth condition or the doubling condition, the theorem is still available for $p = q$. In fact, under the growth condition or the doubling condition for any cube $Q \in \mathcal{Q}(\mu)$ we can find a large integer $j \gg 1$ such that $2^j Q \in \mathcal{Q}(\mu; k, \beta)$.

Remark 2.3. This theorem readily extends to the vector-valued version. Let $1 \le q \le p < \infty$ and $r \in (1, \infty)$. We define the vector-valued Morrey spaces $\mathcal{M}^p_q(l^r, \mu)$ by the set of sequences of μ-measurable functions $\{f_j\}_{j \in \mathbf{N}}$ for which

$$\|f_j \,:\, \mathcal{M}^p_q(l^r, \mu)\| := \sup_{Q \in \mathcal{Q}(\mu)} \mu(2Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q \|f_j \,:\, l^r\|^q \, d\mu \right)^{\frac{1}{q}} < \infty.$$

The theorem can be extended to the vector valued version. Let

$$\|f_j \,:\, \mathcal{M}^p_q(l^r, \mu)\|_d := \sup_{Q \in \mathcal{Q}(\mu;k,\beta)} \mu(Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q \|f_j(y) \,:\, l^r\|^q \, d\mu(y) \right)^{\frac{1}{q}}.$$

Then $C^{-1} \|f_j \,:\, \mathcal{M}^p_q(l^r, \mu)\|_d \le \|f_j \,:\, \mathcal{M}^p_q(l^r, \mu)\| \le C \|f_j \,:\, \mathcal{M}^p_q(l^r, \mu)\|_d$. The same proof as the scalar-valued spaces works for the vector-valued spaces, so in the actual proof we concentrate on the scalar-valued cases.

Proof. Given $k > 1$, we shall prove

$$C^{-1} \|f \,:\, \mathcal{M}^p_q(\mu)\|_d \le \|f \,:\, \mathcal{M}^p_q(k, \mu)\|, \quad \|f \,:\, \mathcal{M}^p_q(\mu)\| \le C \|f \,:\, \mathcal{M}^p_q(\mu)\|_d$$

for large $\beta > 0$. The left inequality is obvious, so let us prove the right inequality. We have only to show that, for every cube $Q \in \mathcal{Q}(\mu)$,

$$\mu(2Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}} \le C \, \| f \, : \, \mathcal{M}^p_q(\mu) \|_d.$$

Let $x \in Q \cap \mathrm{supp}\,(\mu)$ and $Q(x)$ the largest doubling cube centered at x and having sidelength $k^{-j}\ell(Q)$ for some $j \in \mathbf{N}$. Existence of $Q(x)$ can be ensured for μ-almost all $x \in \mathbf{R}^d$. Set

$$\mathcal{Q}_0(j) := \{Q(x) \, : \, \ell(Q(x)) = k^{-j}\ell(Q)\}, \, j \in \mathbf{N}.$$

By Besicovitch's covering lemma we can take $\mathcal{Q}(j) \subset \mathcal{Q}_0(j)$ so that $\sum_{R \in \mathcal{Q}(j)} \chi_R \le 4^d \chi_{2Q}$ and that $x \in \bigcup_{R \in \mathcal{Q}(j)} R$ for μ-almost all $x \in Q$ with $\ell(Q(x)) = k^{-j}\ell(Q)$. Volume argument gives us that $\sharp(\mathcal{Q}(j)) \le 8^d \, k^{jd}$. Since $\left(\int_Q |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}} \le \sum_{j=1}^{\infty} \sum_{R \in \mathcal{Q}(j)} \left(\int_R |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}}$ and $\mu(R) \le \beta^{-j}\mu(2Q)$ for all $R \in \mathcal{Q}(j)$, we have

$$\begin{aligned}
\mu(2Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}} &\le \sum_{j=1}^{\infty} \beta^{j\left(\frac{1}{p}-\frac{1}{q}\right)} \sum_{R \in \mathcal{Q}(j)} \mu(R)^{\frac{1}{p}-\frac{1}{q}} \left(\int_R |f(y)|^q \, d\mu(y) \right)^{\frac{1}{q}} \\
&\le \sum_{j=1}^{\infty} 8^d \, k^{jd} \, \beta^{j\left(\frac{1}{p}-\frac{1}{q}\right)} \| f \, : \, \mathcal{M}^p_q(\mu) \|_d \\
&= \sum_{j=1}^{\infty} 8^d \, \exp\left\{ j \left(d \log k + \left(\frac{1}{p} - \frac{1}{q} \right) \log \beta \right) \right\} \| f \, : \, \mathcal{M}^p_q(\mu) \|_d \le C \, \| f \, : \, \mathcal{M}^p_q(\mu) \|_d,
\end{aligned}$$

where the constant C is finite, provided $\beta > k^{\frac{dpq}{p-q}}$. ∎

3 Equivalent norms of Campanato type

Throughout the rest of this paper we assume that μ satisfy the growth condition (1). We do not assume that μ is doubling. Before we formulate our theorems, let us recall the definition of the RBMO spaces due to Tolsa [10]. Given two cubes $Q \subset R$ with $Q \in \mathcal{Q}(\mu)$, we denote

$$\delta(Q, R) := \int_{\ell(Q)}^{\ell(Q_R)} \frac{\mu(Q(z_Q, l))}{l^n} \, \frac{dl}{l}, \quad K_{Q,R} = 1 + \delta(Q, R),$$

where Q_R denotes the smallest cube concentric to Q containing R. Here and below we abbreviate the $(2, 2^{d+1})$-doubling cube to the doubling cube and $\mathcal{Q}(\mu; 2, 2^{d+1})$ to $\mathcal{Q}(\mu, 2)$. Given $Q \in \mathcal{Q}(\mu)$, we set Q^* as the smallest doubling cube R of the form $R = 2^j Q$ with $j = 0, 1, \ldots$.[2]

Tolsa defined a new BMO for the growth measures, which is suitable for the Calderón-Zygmund theory. We say that $f \in L^1_{loc}(\mu)$ is an element of RBMO if it satisfies

$$\|f\|_* := \sup_{Q \in \mathcal{Q}(\mu)} \frac{1}{\mu\left(\frac{3}{2}Q\right)} \int_Q |f(x) - m_{Q^*}(f)|\, d\mu(x) + \sup_{\substack{Q \subset R \\ Q, R \in \mathcal{Q}(\mu,2)}} \frac{|m_Q(f) - m_R(f)|}{K_{Q,R}} < \infty,$$

where $m_Q(f) := \dfrac{1}{\mu(Q)} \displaystyle\int_Q f(y)\, d\mu(y)$. Further details may be found in [10, Section 2]. The following lemma is due to Tolsa.

Lemma 3.1. [10, Corollary 3.5] *Let $f \in$RBMO.*

1. *There exist positive constants C and C' independent of f so that, for every $\lambda > 0$ and every cube $Q \in \mathcal{Q}(\mu)$,*

$$\mu\{x \in Q \,:\, |f(x) - m_{Q^*}(f)| > \lambda\} \le C\, \mu\left(\frac{3}{2}Q\right) \exp\left(-\frac{C'\lambda}{\|f\|_*}\right).$$

2. *Let $1 \le q < \infty$. Then there exists a constant C independent of f, so that, for every cube $Q \in \mathcal{Q}(\mu)$,*

$$\left(\frac{1}{\mu\left(\frac{3}{2}Q\right)} \int_Q |f(x) - m_{Q^*}(f)|^q\, d\mu(x)\right)^{\frac{1}{q}} \le C\, \|f\|_*.$$

Elementary property of $\delta(\cdot, \cdot)$ Below we list elementary properties of $\delta(\cdot, \cdot)$ used in this paper.

Lemma 3.2. *Let $Q \in \mathcal{Q}(\mu)$. Then the following properties hold :*

(1) *For $\rho > 1$, we have $\delta(Q, \rho Q) \le C_0 \log \rho$.*

(2) $\delta(Q, Q^*) \le C_0\, 2^{n+1} \log 2$.

(3) *Let $k_0 \in \mathbf{N}$ and $\alpha > 0$. Assume, for some $\theta > 0$, $\alpha \le \mu(Q) \le \mu(2^{k_0} Q) \le \theta\, \alpha$. Then $\delta(Q, 2^{k_0} Q) \le 2^n \log 2 \cdot \theta\, C_0\, c_n$, where $c_n := \displaystyle\sum_{k=0}^{\infty} 2^{-nk}$.*

[2] By the growth condition (1) there are a lot of big doubling cubes. Precisely speaking, given a cube $Q \in \mathcal{Q}(\mu)$, we can find $j \in \mathbf{N}$ with $2^j Q \in \mathcal{Q}(\mu, 2)$ (see [10]).

(4) *Given the cubes $P \subset Q \subset R$ with $P \in \mathcal{Q}(\mu)$, then*

$$|\delta(P,R) - (\delta(P,Q) + \delta(Q,R))| \le C,$$

where C is a constant depending only on C_0, n, d.

(5) *Let $Q, R \in \mathcal{Q}(\mu)$. Suppose, for some constant $c_1 > 1$, $Q \subset R$ and $\ell(R) \le c_1\,\ell(Q)$. Then there exists a doubling cube $S \in \mathcal{Q}(\mu, 2)$ such that $Q^*, R^* \subset S$ and $\delta(Q^*, S), \delta(R^*, S) \le C$, where C is a constant depending only on c_1, C_0, n, d.*

Proof. In [9], we have proved (1)–(4). For reader's convenience the full proof is given here. (1) is obvious. To prove (2) we set $Q^* = 2^{k_0}Q_0$. We may assume that $k_0 \ge 1$. The dyadic argument yields that $\delta(Q, 2^{k_0}Q) = \displaystyle\int_{\ell(Q)}^{\ell(2^{k_0}Q)} \frac{\mu(Q(z_Q,l))}{l^n}\,\frac{dl}{l} \le 2^n\,\log 2 \sum_{k=1}^{k_0} \frac{\mu(2^kQ)}{\ell(2^kQ)^n}$. Note that $2^{d+1}\,\mu(2^{k-1}Q) \le \mu(2^kQ)$ for $k = 1, 2, \ldots, k_0$, since $2^{k-1}Q$ is not doubling, which yields, together with the fact that $d \ge n$,

$$\delta(Q, 2^{k_0}Q) \le 2^n\,\log 2\,\frac{\mu(2^{k_0}Q)}{\ell(2^{k_0}Q)^n} \sum_{k=1}^{k_0} (2^{n-d-1})^{k_0-k} \le C_0\,2^{n+1}\,\log 2.$$

We prove (3). It follows by the dyadic argument and the assumption that

$$\delta(Q, 2^{k_0}Q) \le 2^n\,\log 2 \sum_{k=1}^{k_0} \frac{\mu(2^kQ)}{\ell(2^kQ)^n} \le 2^n\,\log 2 \cdot \frac{\theta\alpha}{\ell(Q)^n} \sum_{k=0}^{k_0} 2^{-nk} \le 2^n\,\log 2 \cdot \theta\, C_0\, c_n.$$

Now we prove (4). It suffices to prove that

$$A := |\delta(P_Q, R) - \delta(Q, R)| \le C. \tag{2}$$

We decompose A as

$$\begin{aligned} A \;&=\; \left| \int_{\ell(P_Q)}^{\ell(P_R)} \frac{\mu(Q(z_P,l))}{l^n}\,\frac{dl}{l} - \int_{\ell(Q)}^{\ell(Q_R)} \frac{\mu(Q(z_Q,l))}{l^n}\,\frac{dl}{l} \right| \\ &\le\; \int_{\ell(Q)}^{\ell(P_Q)} \frac{\mu(Q(z_Q,l))}{l^n}\,\frac{dl}{l} + \left| \int_{\ell(P_Q)}^{\min\{\ell(P_R),\ell(Q_R)\}} (\mu(Q(z_P,l)) - \mu(Q(z_Q,l)))\,\frac{dl}{l^{n+1}} \right| \\ &\quad + \int_{\min\{\ell(P_R),\ell(Q_R)\}}^{\max\{\ell(P_R),\ell(Q_R)\}} \left(\frac{\mu(Q(z_P,l))}{l^n} + \frac{\mu(Q(z_Q,l))}{l^n} \right) \frac{dl}{l} \;=:\; A_1 + A_2 + A_3. \end{aligned}$$

By (1) the integrals A_1 and A_3 are easily estimated above by some constant C. So we estimate A_2. Bound A_2 from above by

$$A_2 \le \int_{\ell(P_Q)}^{\infty} \mu(Q(z_P,l)\Delta Q(z_Q,l))\,\frac{dl}{l^{n+1}} = \int_{\ell(P_Q)}^{\infty}\int_{\mathbf{R}^d} \chi_{Q(z_P,l)\Delta Q(z_Q,l)}(y)\,d\mu(y)\,\frac{dl}{l^{n+1}}.$$

A simple geometric observation tells us that $\chi_{Q(z_P,l)\Delta Q(z_Q,l)}(y) = 0$ if

$$l \notin \left[\min\{|y-z_P|_\infty, |y-z_Q|_\infty\},\, \max\{|y-z_P|_\infty, |y-z_Q|_\infty\}\right],$$

where $|y|_\infty := \max\{|y_1|,\dots,|y_d|\}$. This observation and Fubini's theorem yield

$$\begin{aligned} A_2 &\le C\int_{\mathbf{R}^d\setminus P_Q}\left|\frac{1}{|y-z_P|_\infty{}^n} - \frac{1}{|y-z_Q|_\infty{}^n}\right| d\mu(y) \\ &\le C\int_{|y-z_P|_\infty \ge \ell(P_Q)/2} \frac{|z_P-z_Q|_\infty}{|y-z_P|_\infty{}^{n+1}}\,d\mu(y) \le C\,\frac{|z_P-z_Q|_\infty}{\ell(P_Q)} \le C. \end{aligned}$$

This proves (2).

Finally we establish (4). Let $Q^* = 2^jQ$. Then we claim $\delta(R, 2^jR) \le C$. Indeed, by virtue of the fact that $Q \subset R$ we see that if $l \ge \ell(R)$ then $Q(z_R,l) \subset Q(z_q,2l)$. As a consequence we obtain

$$\begin{aligned} \delta(R,2^jR) &= \int_{\ell(R)}^{2^j\ell(R)} \frac{\mu(Q(z_R,l))}{l^n}\,\frac{dl}{l} \\ &\le \int_{\ell(R)}^{2^j\ell(R)} \frac{\mu(Q(z_Q,2l))}{l^n}\,\frac{dl}{l} \le \int_{\ell(Q)}^{c_1\,2^{j+1}\ell(Q)} \frac{\mu(Q(z_Q,l))}{l^n}\,\frac{dl}{l} \le C. \end{aligned}$$

If we put $S := (2^{j+1}R)^*$, then $\delta(R^*,S) \le C$. (1) and (4) finally give us

$$\delta(Q^*,S) \le \delta(Q^*,2^{j+1}R) + \delta(2^{j+1}R,S) + C \le C.$$

This is the desired result. ∎

Scalar-valued Campanato space Having cleared up the definition of RBMO, we will find a relationship between RBMO and the Morrey spaces. With the definition of RBMO in mind, we shall define the Campanato spaces.

Let $f \in L^1_{loc}(\mu)$. We define the Campanato spaces $\mathcal{C}^p_q(k,\mu)$ normed by

$$\begin{aligned} \|f\,:\,\mathcal{C}^p_q(k,\mu)\| \quad &:= \quad \sup_{Q\in\mathcal{Q}(\mu)} \mu(kQ)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f(x)-m_{Q^*}(f)|^q\,d\mu(x)\right)^{\frac{1}{q}} \\ &\quad + \quad \sup_{\substack{Q\subset R\\ Q,R\in\mathcal{Q}(\mu,2)}} \mu(Q)^{\frac{1}{p}}\frac{|m_Q(f)-m_R(f)|}{K_{Q,R}},\ 1\le q\le p\le\infty,\ k>1. \end{aligned}$$

Let $k_1, k_2 > 1$. Then $\mathcal{C}^p_q(k_1,\mu)$ and $\mathcal{C}^p_q(k_2,\mu)$ coincide as a set and their norms are mutually equivalent. Speaking more precisely, we have the norm equivalence

$$\|f \,:\, \mathcal{C}^p_q(k_1,\mu)\| \sim \|f \,:\, \mathcal{C}^p_q(k_2,\mu)\|. \tag{3}$$

To prove (3) we may assume that $k_2 = 2k_1 - 1$ because of the monotonicity of $\mathcal{C}^p_q(k,\mu)$ with respect to k. Then all we have to prove is

$$\mu(k_1Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q |f(x) - m_{Q^*}(f)|^q \, d\mu(x) \right)^{\frac{1}{q}} \le C \, \|f \,:\, \mathcal{C}^p_q(k_2,\mu)\|$$

for fixed cube $Q \in \mathcal{Q}(\mu)$. Divide equally Q into 2^d cubes and collect those in $\mathcal{Q}(\mu)$. Let us name them $Q_1, Q_2, \ldots, Q_N$, $N \le 2^d$. The triangle inequality reduces the matter to showing

$$\mu(k_1Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_{Q_l} |f(x) - m_{Q^*}(f)|^q \, d\mu(x) \right)^{\frac{1}{q}} \le C \, \|f \,:\, \mathcal{C}^p_q(k_2,\mu)\|, \; 1 \le l \le N.$$

Note that $k_2 Q_l \subset k_1 Q$. We apply Lemma 3.2 (5) to obtain an auxiliary doubling cube R which contains $(Q_l)^*, Q^*$ and satisfies $K_{(Q_l)^*,R}, K_{Q^*,R} \le C$. Thus, we obtain

$$\begin{aligned}
&\mu(k_1Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_{Q_l} |f(x) - m_{Q^*}(f)|^q \, d\mu(x) \right)^{\frac{1}{q}} \\
&\quad\le\quad \mu(k_1Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_{Q_l} |f(x) - m_{(Q_l)^*}(f)|^q \, d\mu(x) \right)^{\frac{1}{q}} \\
&\quad\quad+\quad \mu(Q_l)^{\frac{1}{p}} |m_{(Q_l)^*}(f) - m_R(f)| \;+\; \mu(Q_l)^{\frac{1}{p}} |m_R(f) - m_{Q^*}(f)| \\
&\quad\le\quad C \, \|f \,:\, \mathcal{C}^p_q(k_2,\mu)\|.
\end{aligned}$$

As a result (3) is proved.

Since $\mathcal{C}^p_q(k_1,\mu)$ and $\mathcal{C}^p_q(k_2,\mu)$ are isomorphic to each other as Banach spaces, no confusion can occur if we denote $\mathcal{C}^p_q(\mu) = \mathcal{C}^p_q(2,\mu)$.

Note that $\mathcal{C}^\infty_q(\mu) = RBMO$, if $1 \le q < \infty$. This is an immediate consequence of Lemma 3.1. Thus we can say that RBMO is a limit function space of $\mathcal{C}^p_q(\mu)$ as $p \to \infty$ with $q \in [1,\infty)$ fixed.

Next, we observe $\mathcal{Q}(\mu,2)$ can be seen as a net whose order is induced by natural inclusion. With the aid of the following proposition, we shall cope with the ambiguity of constant functions in the semi-norm of the Campanato spaces.

Proposition 3.3. *Let $1 \le q \le p < \infty$. Then the limit $M(f) := \lim_{Q \in \mathcal{Q}(\mu,2)} m_Q(f)$ exists for every $f \in \mathcal{C}^p_q(\mu)$. That is, given $\varepsilon > 0$, we can find a doubling cube $Q \in \mathcal{Q}(\mu,2)$ such that*

$$|m_R(f) - m_Q(f)| \le \varepsilon$$

for all $R \in \mathcal{Q}(\mu, 2)$ engulfing Q. In particular there exists an increasing sequence of concentric doubling cubes $I_0 \subset I_1 \subset \ldots \subset I_k \subset \ldots$ such that

$$\{m_{I_k}(f)\}_{k\in\mathbf{N}_0} \text{ is Cauchy and } \bigcup_k I_k = \mathbf{R}^d. \tag{4}$$

We remark that the condition like (4) appears in [4]. We are mainly interested in the function $f \in \mathcal{C}_q^p(\mu)$ such that $M(f) = 0$.

Proof. Before we come to the proof of Proposition 3.3, we note that

$$\| \, |f| \, : \, \mathcal{C}_1^p(\mu)\| \le C \, \|f \, : \, \mathcal{C}_1^p(\mu)\|. \tag{5}$$

Indeed, we have

$$\begin{aligned}
&\mu\left(\frac{3}{2}Q\right)^{\frac{1}{p}-1} \int_Q ||f(x)| - m_{Q^*}(|f|)| \, d\mu(x) \\
&= \mu\left(\frac{3}{2}Q\right)^{\frac{1}{p}-1} \frac{1}{\mu(Q^*)} \int_Q \left| \int_{Q^*} |f(x)| - |f(y)| \, d\mu(y) \right| d\mu(x) \\
&\le \mu\left(\frac{3}{2}Q\right)^{\frac{1}{p}-1} \frac{1}{\mu(Q^*)} \int_Q int_{Q^*} \, ||f(x)| - |f(y)|| \, d\mu(y) \, d\mu(x) \\
&\le \mu\left(\frac{3}{2}Q\right)^{\frac{1}{p}-1} \frac{1}{\mu(Q^*)} \int_Q \int_{Q^*} |f(x) - f(y)| \, d\mu(y) \, d\mu(x) \\
&\le \mu\left(\frac{3}{2}Q\right)^{\frac{1}{p}-1} \int_Q |f(x) - m_{Q^*}(f)| \, d\mu(x) + \mu(Q^*)^{\frac{1}{p}-1} \int_{Q^*} |m_{Q^*}(f) - f(y)| \, d\mu(y) \\
&\le C \, \|f \, : \, \mathcal{C}_1^p(\mu)\|.
\end{aligned}$$

In the same way we can prove

$$\sup_{\substack{Q\subset R \\ Q,R\in\mathcal{Q}(\mu,2)}} \mu(Q)^{\frac{1}{p}} \frac{|m_Q(|f|) - m_R(|f|)|}{K_{Q,R}} \le C \, \|f \, : \, \mathcal{C}_1^p(\mu)\|.$$

As a consequence (5) is justified.

We now turn to the proof of Proposition 3.3. By the monotonicity of $\mathcal{C}_q^p(\mu)$ with respect to q, we may assume $q = 1$.

Case 1 μ is infinite. Take a sequence of concentric doubling cubes $\{Q_j\}_{j\in\mathbf{N}}$ such that for all $j \in \mathbf{N}$

$$\mu(Q_1) \ge 1, \, \mu(Q_{j+1}) \ge 2\mu(Q_j), \quad \delta(Q_j, Q_{j+1}) \le C$$

for some $C > 0$ depending only on C_0. Then by the definition of $\mathcal{C}_1^p(\mu)$ it holds that

$$|m_{Q_j}(f) - m_{Q_{j+1}}(f)| \le C\, 2^{-\frac{j}{p}} \|f \,:\, \mathcal{C}_1^p(\mu)\|,\ j \in \mathbf{N}.$$

Thus we establish at least the existence of $M(f) := \lim_{j\to\infty} m_{Q_j}(f)$. Let $Q \in \mathcal{Q}(\mu)(\mu, 2)$ which contains Q_j and does not contain Q_{j+1}. Set $Q' = (Q_j{}^Q)^*$. Then by using Lemma 3.2 it is easy to see that $\delta(Q, Q') \le C$ for some absolute constant $C > 0$. Then we have

$$|m_{Q'}(f) - m_Q(f)|, |m_{Q'}(f) - m_{Q_j}(f)| \le C, \quad 2^{-\frac{j}{p}} \|f \,:\, \mathcal{C}_1^p(\mu)\|,$$

which implies

$$|m_Q(f) - M(f)| \le C\, 2^{-\frac{j}{p}} \|f \,:\, \mathcal{C}_1^p(\mu)\|.$$

Thus we finally establish $M(f) = \lim_{Q\in\mathcal{Q}(\mu,2)} m_Q(f)$.

Case 2 μ is finite. In this case, we have only to prove

Claim 3.4. *If μ is finite and $\|f \,:\, \mathcal{C}_1^p(\mu)\| < \infty$, then $f \in L^1(\mu)$.*

In proving Claim 3.4, (5) allows us to assume f is positive.

We take an increasing sequence of concentric doubling cubes $\{Q_j\}_{j\in\mathbf{N}}$ such that $\delta(Q_1, Q_k) \le C$ for all $k \in \mathbf{N}$. Then we have

$$m_{Q_k}(f) \le m_{Q_1}(f) + \mu(Q_1)^{-\frac{1}{p}}(1 + C)\|f \,:\, \mathcal{C}_q^p(\mu)\|.$$

Passage to the limit then gives

$$\int_{\mathbf{R}^d} f\, d\mu \le \mu(\mathbf{R}^d) \left(m_{Q_1}(f) + \mu(Q_1)^{-\frac{1}{p}}(1 + C)\|f \,:\, \mathcal{C}_q^p(\mu)\| \right).$$

This establishes $f \in L^1(\mu)$. ∎

The main theorem in this section is the following.

Theorem 3.5. *Let $1 \le q \le p < \infty$. Assume $f \in \mathcal{C}_q^p(\mu)$ satisfies $M(f) = \lim_{Q\in\mathcal{Q}(\mu,2)} m_Q(f) = 0$. Then*

$$C^{-1} \|f \,:\, \mathcal{C}_q^p(\mu)\| \le \|f \,:\, \mathcal{M}_q^p(\mu)\| \le C \|f \,:\, \mathcal{C}_q^p(\mu)\|$$

for some constant $C > 0$.

The left inequality is obvious. To prove the right inequality we need a lemma.

Lemma 3.6. *Under the assumption of Theorem 3.5, given $R \in \mathcal{Q}(\mu, 2)$, there exists a sequence of increasing doubling cubes $\{R_k\}_{k=1}^K$ such that*

1. *R_k is concentric and $R_1 = R$.*

2. *If μ is finite, then so is K and $R_K = \mathbf{R}^d$.*

3. *For large $K_0 \in \mathbf{N}$, there exists R_{k_0} so that $R_{k_0} \subset I_{K_0} \subset R_{k_0+1}$.*

4. *$\mu(R_k) \geq 2^{k-1}\mu(R)$, $k < K$.*

5. *$\delta(R_k, R_{k+1}) \leq C$, $k < K$.*

Proof. Suppose we have defined R_k. If $\mu(\mathbf{R}^d) \leq 2^k\mu(R)$, then we set $R_{k+1} = \mathbf{R}^d$ and we stop. Suppose otherwise. We define R_{k+1} as the smallest doubling cube of the form $2^l R_k$ with $l \geq 3$ whose μ-measure exceeds $2^k\mu(R)$. By virtue of Lemma 3.2 (3) it is easy to verify that $\{R_k\}_{k=1}^K$ obtained in this way satisfies the property of the lemma. ∎

Let us return to the proof of Theorem 3.5. Let $R \in \mathcal{Q}(\mu)$. We shall estimate

$$\mu(2R)^{\frac{1}{p}-\frac{1}{q}} \left(\int_R |f(x)|^q \, d\mu(x) \right)^{\frac{1}{q}}.$$

The triangle inequality enables us to majorize the above integral by

$$\mu\left(\frac{3}{2}R\right)^{\frac{1}{p}-\frac{1}{q}} \left(\int_R |f(x) - m_{R^*}(f)|^q \, d\mu(x) \right)^{\frac{1}{q}} + \mu(R)^{\frac{1}{p}} |m_{R^*}(f)|.$$

Consequently we can reduce the matters to the estimate of $\mu(R^*)^{\frac{1}{p}}|m_{R^*}(f)|$.

Now we invoke Lemma 3.6 for K_0 taken so that $\mu(R)^{\frac{1}{p}}|m_{I_{K_0}}(f)| \leq \|f : \mathcal{C}_q^p(\mu)\|$. Using the sequence $\{R_k\}_{k=1}^K$, we obtain

$$\begin{aligned} &\mu(R^*)^{\frac{1}{p}} \, |m_{R_k}(f) - m_{R_{k+1}}(f)| \\ &\quad \leq \; C\, 2^{-\frac{k}{p}} \, \mu(R_k)^{\frac{1}{p}} \, \frac{|m_{R_k}(f) - m_{R_{k+1}}(f)|}{1 + \delta(R_k, R_{k+1})} \leq C\, 2^{-\frac{k}{p}} \|f_j : \mathcal{C}_q^p(l^r, \mu)\|. \end{aligned}$$

We also have $\mu(R^*)^{\frac{1}{p}}|m_{R_{k_0}}(f) - m_{I_{K_0}}(f)| \leq C\, 2^{-\frac{k_0}{p}} \|f : \mathcal{C}_q^p(\mu)\|$, since by the properties *3* and *4* of Lemma 3.6 we see that $\delta(R_{k_0}, R_{k_0+1}), \delta(I_{K_0}, R_{k_0+1})$ are majorized by some constants dependent only on C_0. The triangle inequality gives us

$$\begin{aligned} &\mu(R^*)^{\frac{1}{p}} \, |m_{R^*}(f)| \\ &\quad \leq \; \mu(R)^{\frac{1}{p}} \sum_{k=1}^{k_0-1} |m_{R_k}(f) - m_{R_{k+1}}(f)| + \mu(R^*)^{\frac{1}{p}} \left(|m_{R_{k_0}}(f) - m_{I_{K_0}}(f)| + |m_{I_{K_0}}(f)| \right) \\ &\quad \leq \; C \left(\sum_{k=1}^{\infty} 2^{-\frac{k}{p}} \right) \|f : \mathcal{C}_q^p(\mu)\| + \mu(R^*)^{\frac{1}{p}} \, |m_{I_{K_1}}(f)| \leq C \, \|f : \mathcal{C}_q^p(\mu)\|. \end{aligned}$$

The proof of Theorem 3.5 is therefore complete.

Vector-valued extension Finally we consider the vector-valued extensions of Theorem 3.5. Let $\|a_j \,:\, l^r\|$ denote the l^r-norm of $a = \{a_j\}_{j\in\mathbf{N}}$. If possible confusion can occur, then we write $\|\{a_j\}_{j\in\mathbf{N}} \,:\, l^r\|$. For $f \in L^1_{loc}(\mu)$, we define the sharp maximal operator due to Tolsa by

$$M^\sharp f(x) := \sup_{x\in Q\in\mathcal{Q}(\mu)} \frac{1}{\mu\left(\frac{3}{2}Q\right)} \int_Q |f(y) - m_{Q^*}(f)|\, d\mu(y) + \sup_{\substack{x\in Q\subset R \\ Q,R\in\mathcal{Q}(\mu,2)}} \frac{|m_Q(f) - m_R(f)|}{K_{Q,R}}.$$

Lemma 3.1 can be extended to the following vector-valued version.

Lemma 3.7. [9] *Let $f_j \in RBMO$ for $j = 1, 2, \ldots$. For any cube $Q \in \mathcal{Q}(\mu)$ and $q, r \in (1,\infty)$, there exists a constant C independent of f_j such that*

$$\left(\frac{1}{\mu\left(\frac{3}{2}Q\right)} \int_Q \|f_j(x) - m_{Q^*}(f_j) \,:\, l^r\|^q \, d\mu(x)\right)^{\frac{1}{q}} \le C \sup_{x\in\mathbf{R}^d} \left\|M^\sharp f_j(x) \,:\, l^r\right\|. \quad (6)$$

We now define the vector-valued Campanato spaces. Let $1 \le q \le p \le \infty$ and $r \in (1,\infty)$. We say that $\{f_j\}_{j\in\mathbf{N}}$ belongs to the vector-valued Campanato spaces $\mathcal{C}^p_q(l^r,\mu)$ if each f_j is μ-measurable and

$$\begin{aligned} \|f_j \,:\, \mathcal{C}^p_q(l^r,\mu)\| \quad &:= \quad \sup_{Q\in\mathcal{Q}(\mu)} \mu(2Q)^{\frac{1}{p}-\frac{1}{q}} \left(\int_Q \|f_j(x) - m_{Q^*}(f_j) \,:\, l^r\|^q \, d\mu(x)\right)^{\frac{1}{q}} \\ &+ \quad \sup_{\substack{Q\subset R \\ Q,R\in\mathcal{Q}(\mu,2)}} \mu(Q)^{\frac{1}{p}} \frac{\|m_Q(f_j) - m_R(f_j) \,:\, l^r\|}{K_{Q,R}} < \infty. \end{aligned}$$

As for the vector-valued spaces, the norm equivalence of the Campanato type still holds.

Theorem 3.8. *Let $1 \le q \le p < \infty$ and let $\{f_j\}_{j\in\mathbf{N}}$ be a sequence in $\mathcal{C}^p_q(\mu)$. Assume that there exists an increasing sequence of concentric doubling cubes $I_0 \subset I_1 \subset \ldots \subset I_k \subset \ldots$ such that*

$$\lim_{k\to\infty} m_{I_k}(f_j) = 0 \text{ for all } j \text{ and } \bigcup_k I_k = \mathbf{R}^d.$$

Then there exists a constant $C > 0$ independent of $\{f_j\}_{j\in\mathbf{N}}$ such that

$$C^{-1} \|f_j \,:\, \mathcal{C}^p_q(l^r,\mu)\| \le \|f_j \,:\, \mathcal{M}^p_q(l^r,\mu)\| \le C \|f_j \,:\, \mathcal{C}^p_q(l^r,\mu)\|.$$

Using Lemma 3.7, we can say more about $\mathcal{C}^\infty_q(l^r,\mu)$, which gives us a partial clue to the definition of the vector-valued RBMO spaces. Speaking precisely, we obtain the following proposition.

Proposition 3.9. *Let $\{f_j\}_{j\in\mathbf{N}}$ be a sequence of $L^1_{loc}(\mu)$ functions. Then*

$$\sup_{\substack{Q\subset R\\ Q,R\in\mathcal{Q}(\mu,2)}} \frac{\|m_Q(f_j)-m_R(f_j)\ :\ l^r\|}{K_{Q,R}} \le c\ \sup_{x\in\mathbf{R}^d}\|M^\sharp f_j(x)\ :\ l^r\|. \tag{7}$$

In particular, we have

$$\|f_j\ :\ \mathcal{C}^\infty_q(l^r,\mu)\| \le c\ \sup_{x\in\mathbf{R}^d}\|M^\sharp f_j(x)\ :\ l^r\|. \tag{8}$$

Proof. Fix $Q\subset R$ such that $Q\in\mathcal{Q}(\mu)$. Then $\dfrac{|m_Q(f_j)-m_R(f_j)|}{K_{Q,R}} \le c\,M^\sharp f_j(x)$ for all $x\in Q$. By taking the l^r-norm of both sides we obtain

$$\frac{\|m_Q(f_j)-m_R(f_j)\ :\ l^r\|}{K_{Q,R}} \le c\sup_{x\in Q}\|M^\sharp f_j(x)\ :\ l^r\| \le c\ \sup_{x\in\mathbf{R}^d}\|M^\sharp f_j(x)\ :\ l^r\|.$$

Now since Q and R are taken arbitrarily, (7) is proved. (8) can be obtained with the help of (6) and (7). ∎

Before we conclude this section, a remark may be in order.

Remark 3.10. Let $0<\alpha<n$. For $Q,R\in\mathcal{Q}(\mu)$ with $Q\subset R$ we define

$$K^{(\alpha)}_{Q,R} = 1+\sum_{k=1}^{N_{Q,R}}\left(\frac{\mu(2^kQ)}{\ell(2^kQ)^n}\right)^{\frac{n-\alpha}{n}},$$

where $N_{Q,R}$ is the least integer j with $2^jQ\supset R$. For the definition of this constant we refer to [3]. Theorems in this section still hold, if we replace $K_{Q,R}$ by $K^{(\alpha)}_{Q,R}$ whenever $1\le q\le p<\infty$.

Acknowledgement

The authors thank Prof. E. Nakai for discussing this paper with them.

References

[1] S. Campanato, Proprietà di una famiglia di spazi funzionali. (Italian) Ann. Scuola Norm. Sup. Pisa (3) 18 1964 137–160.

[2] S. Campanato, Proprietà di inclusione per spazi di Morrey. (Italian) Ricerche Mat. 12 1963 67–86.

[3] W. Chen and E. Sawyer, A note on commutators of fractional integrals with RBMO(μ) functions. Illinois J. Math. **46** (2002), no. 4, 1287-1298.

[4] N. Fujii, A proof of the Fefferman-Stein-Strömberg inequality for the sharp maximal functions, Proc. Amer. Soc. **106** (1991), no.2, 371–377.

[5] C. Morrey, On solutions of quasi-linear elliptic partial differential equations, Trans. Amer. Math. Soc. **43** (1938), 126–166.

[6] E. Nakai, The Campanato, Morrey and Hölder spaces on spaces of homogeneous type, in preparation.

[7] Y. Sawano and H. Tanaka, Morrey spaces for non-doubling measures, Acta Math. Sinica **21** no.6, 1535–1544.

[8] Y. Sawano and H. Tanaka, Sharp maximal inequalities and commutators on Morrey spaces with non-doubling measures, to appear in Taiwanese Math.J..

[9] Y. Sawano and H. Tanaka, The John-Nirenberg type inequality for non-doubling measures, *preprint*.

[10] X. Tolsa, BMO, H^1, and Calderón-Zygmund operators for non doubling measures, Math. Ann. **319** (2001), 89-149.

[11] X. Tolsa, The space H^1 for nondoubling measures in terms of a grand maximal operator, Trans. Amer. Math. Soc., **355** (2003), 315–348.